

Classification of the BPS states in Bagger-Lambert Theory

IMTAK JEON,[†] JONGWOOK KIM,[†] NAKWOO KIM,^{*}

SANG-WOO KIM[‡] AND JEONG-HYUCK PARK[†]

[†]Department of Physics, Sogang University, Seoul 121-742, Korea

^{*}Department of Physics and Research Institute of Basic Science
Kyung Hee University, Seoul 130-701, Korea

[‡]Center for Quantum Spacetime, Sogang University, Seoul 121-742, Korea

Abstract

We classify, in a group theoretical manner, the BPS configurations in the multiple M2-brane theory recently proposed by Bagger and Lambert. We present three types of BPS equations preserving various fractions of supersymmetries: in the first type we have constant fields and the interactions are purely algebraic in nature; in the second type the equations are invariant under spatial rotation $SO(2)$, and the fields can be time-dependent; in the third class the equations are invariant under boost $SO(1,1)$ and provide the eleven-dimensional generalizations of the Nahm equations. The BPS equations for different number of supersymmetries exhibit the division algebra structures: octonion, quaternions or complex.

Contents

1	Introduction	2
2	Preliminaries	3
2.1	Supersymmetry projection matrix - general	4
2.2	$SO(1, 2)$ invariant projection matrices	7
2.3	$SO(2)$ invariant projection matrices	7
2.4	$SO(1, 1)$ invariant projection matrices	8
3	Classification of the BPS equations	9
3.1	$SO(1, 2)$ invariant BPS equations	9
3.1.1	$N = 2$ $SO(1, 2) \times SO(7)$ invariant BPS equations - <i>octonion</i>	10
3.1.2	$N = 4$ $SO(1, 2) \times SO(6) \times SO(2)$ invariant BPS equations - <i>complex</i>	11
3.1.3	$N = 6$ $SO(1, 2) \times SO(5) \times SO(3)$ invariant BPS equations - <i>quaternion</i>	12
3.1.4	$N = 8$ $SO(1, 2) \times SO(4) \times SO(4)$ invariant BPS equations	12
3.1.5	$N = 10$ $SO(1, 2) \times SO(3) \times SO(5)$ invariant BPS equations	13
3.1.6	$N = 12$ $SO(1, 2) \times SO(2) \times SO(6)$ invariant BPS equations	13
3.2	$SO(2)$ invariant BPS equations	14
3.2.1	$N = 2$ $SO(2) \times SU(4)$ invariant BPS equations	15
3.2.2	$N = 4$ $SO(2) \times SU(2) \times SO(4)$ invariant BPS equations	15
3.2.3	$N = 6$ $SO(2) \times SO(2) \times SU(3)$ invariant BPS equations	15
3.2.4	$N = 8$ $SO(2) \times SO(2) \times SO(6)$ invariant BPS equations	16
3.3	$SO(1, 1)$ invariant BPS equations	16
3.3.1	$(N_+, N_-) = (1, 0)$ $SO(1, 1) \times SO(7)$ invariant BPS equations - <i>octonion</i>	17
3.3.2	$(N_+, N_-) = (2, 0)$ $SO(1, 1) \times SO(2) \times SO(6)$ invariant BPS equations - <i>complex</i>	17
3.3.3	$(N_+, N_-) = (3, 0)$ $SO(1, 1) \times SO(3) \times SO(5)$ invariant BPS equations - <i>quaternion</i>	17
3.3.4	$(N_+, N_-) = (4, 0)$ $SO(1, 1) \times SO(4) \times SO(4)$ invariant BPS equations	17
3.3.5	$(N_+, N_-) = (5, 0)$ $SO(1, 1) \times SO(5) \times SO(3)$ invariant BPS equations	18
3.3.6	$(N_+, N_-) = (6, 0)$ $SO(1, 1) \times SO(6) \times SO(2)$ invariant BPS equations	18
3.3.7	$(N_+, N_-) = (7, 0)$ $SO(1, 1) \times SO(7)$ invariant BPS equations	18
3.3.8	$(N_+, N_-) = (1, 1)$ $SO(1, 1) \times SO(6)$ invariant BPS equations	19
3.3.9	$(N_+, N_-) = (2, 2)$ $SO(1, 1) \times SO(2) \times SO(2) \times SO(4)$ invariant BPS equations	19
3.3.10	$(N_+, N_-) = (3, 3)$ $SO(1, 1) \times SO(3) \times SO(3) \times SO(2)$ invariant BPS equations	19
3.3.11	$(N_+, N_-) = (4, 4)$ $SO(1, 1) \times SO(4) \times SO(4)$ invariant BPS equations	19
4	Discussion	20
A	Gamma matrices and octonions	22
B	$SO(2)$ invariant projection matrix	24

1 Introduction

In a series of recent papers [1], Bagger and Lambert (BL) have constructed a three-dimensional, interacting superconformal gauge theory of multiple M2-branes. The action is maximally supersymmetric with 16 ordinary supersymmetries, and it has been verified that the theory is indeed superconformal with 16 conformal supercharges in [2]. In the quest for the final form of the theory, as usual, it was supersymmetry that provided crucial guiding lights. The work was initiated as an attempt to incorporate Basu and Harvey's generalized Nahm equation -which was a proposal to describe M2-branes ending on an M5-brane [3]- in the full supersymmetric M2-brane action. Their analysis revealed a novel algebraic structure, namely the 3-algebra, which is also investigated independently by Gustavsson [4]. Since the discovery, the multiple M2-brane theory of Bagger and Lambert has attracted an enormous degree of attention [5–35]. One might expect that, given this genuine superconformal field theory, $\mathcal{M}$ -theory is now about to unveil its mysterious and fundamental features.

In the present paper, we set out to classify the BPS states, or the BPS equations of the BL theory using a group theoretical consideration. Apparently the theory of our interest has the Lorentz group $SO(1, 2)$ and the R-symmetry group $SO(8)$. Instead of providing the full and thorough survey of possible BPS equations, we focus mainly on two different types of BPS equations with different number of supersymmetries, and classify them completely. The first class is completely Lorentz invariant, and the other is invariant under the spatial rotation.

In the first type, the BPS equations are given purely in terms of the three-algebra commutators and independent of the three-dimensional worldvolume coordinates. Thus the corresponding nontrivial configurations possess infinite energy, typically corresponding to BPS objects of infinite size. Previously known analogous algebraic solutions include the longitudinal M5-brane in $\mathcal{M}$ -theory matrix model which is realized in terms of Heisenberg algebra or large N matrices [36].

In the other type the equations are $SO(2)$ rotation invariant, and the fields can be *time-dependent*. A technical reason why we focus on the two classes is that in these cases, fully utilizing the $SO(8)$ triality we are able to classify the BPS equations completely.

In addition to the two classes, there is another possibility to obtain third type of BPS equations via simple tensor product. Namely one can obtain various generalizations of the Nahm equations which are invariant under the boost $SO(1, 1) \subset SO(1, 2)$. Our BPS equations manifest the division algebra structures:

octonion, quaternion or complex. In the paper we will mainly focus on the BPS equations themselves. Our results hold for both the finite and infinite dimensional three-algebras. Note however that the Lorentz invariant BPS equations can have nontrivial solutions only for infinite dimensional three-algebras. The specific solutions and the physical interpretation will be presented in a separate publication [37].

The organization of the present paper is as follows. Sec.2 is for preliminaries. We first discuss the general features of the ‘supersymmetric projection matrices’ and review how to derive the corresponding BPS equations for a given projection matrix. We also explain the relevant symmetries. Then we classify the projection matrices for the $SO(1, 2)$, $SO(2)$ and $SO(1, 1)$ invariant equations. Sec.3 contains our main results of the BPS equations. Sec.3.1 classifies the $SO(1, 2)$ invariant BPS equations preserving two, four, six, eight, ten and twelve supersymmetries.¹ Sec.3.2 classifies the $SO(2)$ invariant BPS equations preserving two, four, six and eight supersymmetries. In Sec.3.3 we discuss the $SO(1, 1)$ invariant BPS equations which generalize the Nahm equations. The final section, Sec.4 contains our results and discussions. In Appendix we review the $SO(8)$ triality and its relation to octonions.

Note added: While this paper is being finished, Ref. [38] appears in ArXiv which partially overlaps with our work, as it discusses the BPS equations of the form: $D_y X_I = \frac{1}{3!} \mathcal{C}_{IJKL} [X^J, X^K, X^L]$. In the present paper, we explicitly spell the coefficients $\mathcal{C}_{IJKL}$ and classify various BPS equations.

2 Preliminaries

The multiple M2-brane theory has 8 real scalar fields $X^I, I = 1, 2, \dots, 8$ and a 16 component Majorana spinor Ψ . The supersymmetry transformation of the fermions in the Bagger-Lambert theory assumes the form:

$$\delta\Psi = \left(F_{\mu I} \Gamma^{\mu I} - \frac{1}{6} F_{IJK} \Gamma^{IJK} \right) \varepsilon, \quad (2.1)$$

where all the variables are three-algebra valued and we set

$$F_{\mu I} \equiv D_{\mu} X_I, \quad F_{IJK} \equiv [X_I, X_J, X_K]. \quad (2.2)$$

¹Note that in the present paper we focus on the sixteen ordinary supersymmetries and not the sixteen conformal supersymmetries. For the BPS equations preserving conformal supersymmetries in super Yang-Mills we refer the readers to Ref. [39].

The bracket $[X_I, X_J, X_K]$ denotes the three-algebra product which is trilinear and totally antisymmetric. Note also that in contrast to the original convention [1] we let $I = 1, 2, \dots, 8$ and take $\mu \equiv 0, 9, 10$ directions as for the M2-brane worldvolume for convenience to present the BPS equations later,

$$x^0 \equiv t, \quad x^9 \equiv x, \quad x^{10} \equiv y. \quad (2.3)$$

The supersymmetry parameter is real and subject to the $\text{SO}(1, 2)$ projection condition:

$$\Gamma^{txy} \varepsilon = \varepsilon, \quad (2.4)$$

which is consistent with the opposite projection property, $\Gamma^{txy} \Psi = -\Psi$. Since the product of all the eleven-dimensional gamma matrices leads to the 32×32 identity matrix $\Gamma^{txy123\dots 8} = 1$, the above $\text{SO}(1, 2)$ projection condition coincides with the chirality condition of $\text{SO}(8)$,

$$\Gamma^{123\dots 8} \varepsilon = \varepsilon. \quad (2.5)$$

2.1 Supersymmetry projection matrix - general

In general for supersymmetric theories, the supersymmetry projection matrix Ω can be defined in terms of the commuting, real, orthonormal supersymmetry parameters $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_N$,

$$\Omega := \sum_{i=1}^N \varepsilon_i \varepsilon_i^\dagger, \quad \varepsilon_i^\dagger \varepsilon_j = \delta_{ij}, \quad (2.6)$$

satisfying $\Omega^\dagger = \Omega^2 = \Omega$. Here N denotes the number of the preserved supersymmetries,

$$N = \text{Tr} \Omega. \quad (2.7)$$

Naturally the eigenvalues of the projection matrices are either zero or one.

When the supersymmetry transformation of fermions takes the form $\delta \Psi = \mathcal{F} \varepsilon$ where $\mathcal{F}$ denotes a bosonic quantity contracted with gamma matrices as in (2.1), the general strategy to obtain the BPS equations is as follows [40]:

1. *Expand the projection matrix Ω in terms of the gamma matrix product basis.*
2. *Perform the matrix product $\mathcal{F} \Omega$ and reexpress it in terms of the gamma matrix product basis.*

3. Read off the BPS equations from the coefficients of the linearly independent terms.

For example in the Euclidean four-dimensional minimal super Yang-Mills theory, we have two choices for the projection matrix $\Omega = \frac{1}{2}(1 \pm \gamma^{1234})$, while $\mathcal{F} = F_{ij}\gamma^{ij}$. Consequently, noting $\gamma^{12}\Omega = \mp\gamma^{34}\Omega$ etc., we get $F_{ij}\gamma^{ij}\Omega = 2(F \mp \star F)_{i4}\gamma^{i4}\Omega$ such that the corresponding BPS equations are the well-known self-dual or anti-self-dual equations $F = \pm\star F$. In this way, the complete classifications of the BPS equations in six and eight-dimensional super Yang-Mills as well as the pp-wave M-theory matrix model [41] have been carried out [40, 42, 43].

The present paper concerns the BPS equations of the Bagger-Lambert theory. Since the eleven-dimensional spacetime admits Majorana spinors we can set all the gamma matrices and the spinors to be real. In particular, the spatial gamma matrices are symmetric while the temporal gamma matrix is anti-symmetric. Consequently, also from (2.5), the projection matrices of the Bagger-Lambert theory must satisfy

$$\Omega = \Omega^T = \Omega^*, \quad \Omega = \Omega^2, \quad \Omega = \mathcal{P}\Omega = \Omega\mathcal{P}, \quad (2.8)$$

where $\mathcal{P}$ is the $\text{SO}(8)$ chiral projection matrix,

$$\mathcal{P} := \frac{1}{2}(1 + \Gamma^{123\cdots 8}). \quad (2.9)$$

The most general form of such projection matrices reads

$$\Omega = [c + \Upsilon_4 + \Gamma^x(c' + \Upsilon'_4) + \Gamma^y(c'' + \Upsilon''_4) + \Gamma^{xy}\Upsilon_2] \mathcal{P}, \quad (2.10)$$

where c, c', c'' are constants, $\Upsilon_4, \Upsilon'_4, \Upsilon''_4$ are foursome productions of the $\text{SO}(8)$ gamma matrices Γ^{IJKL} contracted with self-dual four-forms, and Υ_2 is a twosome production of the $\text{SO}(8)$ gamma matrices Γ^{IJ} contracted with a two-form. All together, *a priori*, there are $3 + 3 \times \frac{1}{2} \binom{8}{4} + \binom{8}{2} = 136$ real parameters which must be determined by requiring the remaining condition $\Omega^2 = \Omega$. The symmetry group $\text{SO}(1, 2) \times \text{SO}(8)$ in the Bagger-Lambert theory may reduce the number of the free parameters, but is not big enough to transform all the free parameters, the two-form and the four-forms, into ‘canonical’ forms. Note that the $\text{SO}(8)$ rotation may take only one of $\{\Upsilon_4, \Upsilon'_4, \Upsilon''_4, \Upsilon_2\}$ into a canonical form. In our choice, the canonical form of a two-form reads

$$\Upsilon_2 = a_1\Gamma^{12} + a_2\Gamma^{34} + a_3\Gamma^{56} + a_4\Gamma^{78}, \quad (2.11)$$

while the canonical form of a self-dual four-form reads

$$\Upsilon_4 = b_1 \mathcal{E}_1 + b_2 \mathcal{E}_2 + b_3 \mathcal{E}_3 + b_4 \mathcal{E}_4 + b_5 \mathcal{E}_5 + b_6 \mathcal{E}_6 + b_7 \mathcal{E}_7, \quad (2.12)$$

where we set

$$\begin{aligned} \mathcal{E}_1 &= \Gamma_{8127} \mathcal{P}, & \mathcal{E}_2 &= \Gamma_{8163} \mathcal{P}, & \mathcal{E}_3 &= \Gamma_{8246} \mathcal{P}, & \mathcal{E}_4 &= \Gamma_{8347} \mathcal{P}, \\ \mathcal{E}_5 &= \Gamma_{8567} \mathcal{P}, & \mathcal{E}_6 &= \Gamma_{8253} \mathcal{P}, & \mathcal{E}_7 &= \Gamma_{8154} \mathcal{P}. \end{aligned} \quad (2.13)$$

The former is well known, while the latter is less familiar and we review it in Appendix A. In (2.13) the subscript spatial indices of the gamma matrices are organized such that the three indices after the common 8 are identical to those of the totally anti-symmetric octonionic structure constants [40, 44]:

$$\begin{aligned} e_i e_j &= -\delta_{ij} + c_{ijk} e_k, & i, j, k &= 1, 2, \dots, 7, \\ 1 &= c_{127} = c_{163} = c_{246} = c_{347} = c_{567} = c_{253} = c_{154}, & \text{others zero.} \end{aligned} \quad (2.14)$$

We say Ω is invariant under $\text{SO}(2)$ rotation invariant on xy -plane if $[\Gamma_{xy}, \Omega] = 0$. When this holds, for a finite angle ϕ and rotation $G = e^{\phi \Gamma_{xy}}$, from the equivalence

$$\mathcal{F}\Omega = 0 \quad \Longleftrightarrow \quad G\mathcal{F}\Omega G^{-1} = G\mathcal{F}G^{-1}\Omega = 0, \quad (2.15)$$

we note that the corresponding BPS equations are, as a set, invariant under the rotation. Naturally this generalizes to an arbitrary subgroup of $\text{SO}(1, 2) \times \text{SO}(8)$.

In the present paper instead of attempting to solve for the most general projection matrices, we restrict to the cases where Ω assumes the canonical form. Namely we focus on two types of the BPS equations and classify the corresponding BPS equations completely: one is the $\text{SO}(1, 2)$ *invariant cases* i.e.

$$\Omega = (c + \Upsilon_4) \mathcal{P}, \quad (2.16)$$

and the other is the $\text{SO}(2)^5 \equiv \text{SO}(2) \times \text{SO}(2) \times \text{SO}(2) \times \text{SO}(2) \times \text{SO}(2)$ *invariant cases* i.e.

$$\Omega = (\text{constant} + \text{twosome products of } \{\Gamma^{xy}, \Gamma^{12}, \Gamma^{34}, \Gamma^{56}, \Gamma^{78}\}) \mathcal{P}. \quad (2.17)$$

Here $\text{SO}(1, 2)$ and $\text{SO}(2)$ correspond to the M2 worldvolume Lorenz symmetry and the Cartan subgroup of the symmetry group $\text{SO}(1, 2) \times \text{SO}(8)$ respectively. In addition, the former will easily generate various

M-theoretic generalizations of the Nahm equations which are invariant under $\text{SO}(1, 1) \subset \text{SO}(1, 2)$, as the corresponding projection matrices are of the form:

$$\Omega = (1 \pm \Gamma^{tx}) (c + \Upsilon_4) \mathcal{P}. \quad (2.18)$$

2.2 $\text{SO}(1, 2)$ invariant projection matrices

The basic building blocks of all the possible $\text{SO}(1, 2)$ invariant projection matrices are the following $N = 2$ projection matrices [40]:

$$\Omega = \frac{1}{8} (\mathcal{P} + \alpha_1 \alpha_2 \mathcal{E}_1 + \alpha_1 \alpha_3 \mathcal{E}_2 + \alpha_3 \mathcal{E}_3 + \alpha_2 \mathcal{E}_4 + \alpha_1 \mathcal{E}_5 + \alpha_1 \alpha_2 \alpha_3 \mathcal{E}_6 + \alpha_2 \alpha_3 \mathcal{E}_7), \quad (2.19)$$

where $\alpha_1, \alpha_2, \alpha_3$ are three independent signs,

$$\alpha_1^2 = \alpha_2^2 = \alpha_3^2 = 1. \quad (2.20)$$

Three independent sign choices lead to eight possible combinations, hence eight $N = 2$ projection matrices. They are orthogonal to each other and complete, as summing all of them gives an identity. Namely they form an orthogonal basis for the $\text{SO}(1, 2)$ invariant projection matrices. General $N = 2k$ projection matrices can be straightforwardly obtained as a k sum of the above eight $N = 2$ projection matrices. Furthermore, from the $\text{SO}(8)$ triality, the $8!/[k!(8-k)!]$ possibilities for the k sum are all equivalent to each other. The corresponding $N = 2k$ BPS equations are $\text{SO}(1, 2) \times \text{SO}(8-k) \times \text{SO}(k)$ invariant.

2.3 $\text{SO}(2)$ invariant projection matrices

The basic building blocks of all the possible $\text{SO}(2)$ invariant projection matrices are the following $N = 2$ projection matrices (see Appendix B for derivation):

$$\begin{aligned} \Omega &= \frac{1}{8} [1 + \Gamma^{xy} (\beta_1 \Gamma^{12} + \beta_2 \Gamma^{34} + \beta_3 \Gamma^{56} + \beta_1 \beta_2 \beta_3 \Gamma^{78}) - \beta_1 \beta_2 \Gamma^{1234} - \beta_3 \beta_1 \Gamma^{1256} - \beta_2 \beta_3 \Gamma^{1278}] \mathcal{P} \\ &= \frac{1}{8} (1 + \beta_1 \Gamma^{xy12}) (1 + \beta_2 \Gamma^{xy34}) (1 + \beta_3 \Gamma^{xy56}) \mathcal{P}, \end{aligned} \quad (2.21)$$

where $\beta_1, \beta_2, \beta_3$ denote three independent signs,

$$\beta_1^2 = \beta_2^2 = \beta_3^2 = 1. \quad (2.22)$$

Eight possible $N = 2$ projection matrices form an orthogonal basis for the $\text{SO}(2)$ invariant projection matrices. General $N = 2k$ projection matrices can be straightforwardly obtained as a k sum of the above eight $N = 2$ projection matrices. However, if the sum contains a pair of two opposite overall sign factors *e.g.* $(+++)$ and $(---)$, the corresponding BPS configurations become $\text{SO}(1, 2)$ invariant as $F_{\mu I} = 0$ and the BPS equations reduce to those of $\text{SO}(1, 2)$ invariant BPS equations. Excluding these cases, up to $\text{SO}(8)$ rotations, there are five inequivalent $\text{SO}(2)$ invariant projection matrices as follows.

- $N = 2$ $\text{SO}(2) \times \text{SU}(4)$ invariant projection matrix, with the choice of $(\beta_1, \beta_2, \beta_3) = (+++)$,

$$\Omega = \frac{1}{8} [1 + \Gamma^{xy} (\Gamma^{12} + \Gamma^{34} + \Gamma^{56} + \Gamma^{78}) - \Gamma^{1234} - \Gamma^{1256} - \Gamma^{1278}] \mathcal{P}. \quad (2.23)$$

- $N = 4$ $\text{SO}(2) \times \text{SU}(2) \times \text{SO}(4)$ invariant projection matrix, with $(+++), (++-)$,

$$\Omega = \frac{1}{4} [1 + \Gamma^{xy} (\Gamma^{12} + \Gamma^{34}) - \Gamma^{1234}] \mathcal{P}. \quad (2.24)$$

- $N = 6$ $\text{SO}(2) \times \text{SO}(2) \times \text{SU}(3)$ invariant projection matrix, with $(+++), (++-), (+-+)$,

$$\Omega = \frac{1}{8} [3 + \Gamma^{xy} (3\Gamma^{12} + \Gamma^{34} + \Gamma^{56} - \Gamma^{78}) - \Gamma^{1234} - \Gamma^{1256} + \Gamma^{1278}] \mathcal{P}. \quad (2.25)$$

- $N = 8$ $\text{SO}(2) \times \text{SO}(2) \times \text{SO}(6)$ invariant projection matrix, with $(+++), (++-), (+-+), (+--)$,

$$\Omega = \frac{1}{2} (1 + \Gamma^{xy12}) \mathcal{P}. \quad (2.26)$$

- $N = 8$ $\text{SO}(2) \times \text{SU}(4)$ invariant projection matrix, with $(+++), (++-), (+-+), (-++)$,

$$\Omega = \frac{1}{4} [2 + \Gamma^{xy} (\Gamma^{12} + \Gamma^{34} + \Gamma^{56} - \Gamma^{78})] \mathcal{P}. \quad (2.27)$$

2.4 $\text{SO}(1, 1)$ invariant projection matrices

For $\text{SO}(1, 1)$ invariant projection matrices, we have the following $N = 1$ projection matrices:

$$\Omega = \frac{1}{16} (1 + \alpha_0 \Gamma^{tx}) (\mathcal{P} + \alpha_1 \alpha_2 \mathcal{E}_1 + \alpha_1 \alpha_3 \mathcal{E}_2 + \alpha_3 \mathcal{E}_3 + \alpha_2 \mathcal{E}_4 + \alpha_1 \mathcal{E}_5 + \alpha_1 \alpha_2 \alpha_3 \mathcal{E}_6 + \alpha_2 \alpha_3 \mathcal{E}_7), \quad (2.28)$$

where $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ are four independent signs,

$$\alpha_0^2 = \alpha_1^2 = \alpha_2^2 = \alpha_3^2 = 1. \quad (2.29)$$

Sixteen possible $N = 1$ projection matrices form an orthogonal basis for the $\text{SO}(1, 1)$ invariant projection matrices. Generic $N = k$ $\text{SO}(1, 1)$ invariant projection matrices may be obtained straightforwardly as a k sum of the above sixteen $N = 1$ projection matrices. For each sum, we may decompose

$$N = N_+ + N_-, \quad N_+ = n_+ + n, \quad N_- = n_- + n, \quad (2.30)$$

such that $N_{\pm}$ denotes the number of $N = 1$ projection matrices in the sum whose α_0 values are ± 1 , and n counts the number of $N = 1$ projection matrix pairs which have the same $\alpha_1, \alpha_2, \alpha_3$ values and opposite α_0 signs. There are $8!/[n_+!n_-!n!(8-n_+-n_-+n)!]$ possibilities for the sum which are all equivalent to another, thanks to the $\text{SO}(8)$ triality. Furthermore, if n is nontrivial $n \neq 0$, then the BPS configurations become $\text{SO}(1, 2)$ invariant as $F_{\mu I} = 0$ and the number of the preserved supersymmetries is automatically increased from $n_+ + n_- + 2n$ to $2(n_+ + n_- + n)$. In this case the BPS equations reduce to those of $\text{SO}(1, 2)$ invariant BPS equations. Genuinely $\text{SO}(1, 1)$ invariant BPS equations appear only when $n = 0$. The corresponding (N_+, N_-) BPS equations are then $\text{SO}(1, 1) \times \text{SO}(N_+) \times \text{SO}(N_-) \times \text{SO}(8 - N_+ - N_-)$ invariant with the natural restriction $N_+ + N_- \leq 8$.

3 Classification of the BPS equations

3.1 $\text{SO}(1, 2)$ invariant BPS equations

The generic $N = 2$ $\text{SO}(1, 2)$ invariant projection matrix (2.19) leads to the following $N = 2$ $\text{SO}(1, 2) \times \text{SO}(7)$ invariant BPS equations which involve three free sign factors $\alpha_1^2 = \alpha_2^2 = \alpha_3^2 = 1$:

$$F_{\mu I} = 0, \quad \mu = t, x, y, \quad I = 1, 2, \dots, 8, \quad (3.1)$$

and

$$\begin{aligned}
& \alpha_1 \alpha_2 F_{278} + \alpha_2 \alpha_3 F_{548} + \alpha_3 \alpha_1 F_{638} + \alpha_1 F_{234} + \alpha_2 F_{256} + \alpha_3 F_{357} + \alpha_1 \alpha_2 \alpha_3 F_{476} = 0, \\
& \alpha_1 \alpha_2 F_{718} + \alpha_2 \alpha_3 F_{376} + \alpha_3 \alpha_1 F_{475} + \alpha_1 F_{143} + \alpha_2 F_{165} + \alpha_3 F_{468} + \alpha_1 \alpha_2 \alpha_3 F_{538} = 0, \\
& \alpha_1 \alpha_2 F_{456} + \alpha_2 \alpha_3 F_{267} + \alpha_3 \alpha_1 F_{168} + \alpha_1 F_{124} + \alpha_2 F_{478} + \alpha_3 F_{517} + \alpha_1 \alpha_2 \alpha_3 F_{258} = 0, \\
& \alpha_1 \alpha_2 F_{536} + \alpha_2 \alpha_3 F_{158} + \alpha_3 \alpha_1 F_{257} + \alpha_1 F_{132} + \alpha_2 F_{738} + \alpha_3 F_{628} + \alpha_1 \alpha_2 \alpha_3 F_{167} = 0, \\
& \alpha_1 \alpha_2 F_{346} + \alpha_2 \alpha_3 F_{418} + \alpha_3 \alpha_1 F_{427} + \alpha_1 F_{678} + \alpha_2 F_{126} + \alpha_3 F_{137} + \alpha_1 \alpha_2 \alpha_3 F_{328} = 0, \\
& \alpha_1 \alpha_2 F_{354} + \alpha_2 \alpha_3 F_{273} + \alpha_3 \alpha_1 F_{318} + \alpha_1 F_{758} + \alpha_2 F_{152} + \alpha_3 F_{248} + \alpha_1 \alpha_2 \alpha_3 F_{174} = 0, \\
& \alpha_1 \alpha_2 F_{128} + \alpha_2 \alpha_3 F_{236} + \alpha_3 \alpha_1 F_{245} + \alpha_1 F_{568} + \alpha_2 F_{348} + \alpha_3 F_{153} + \alpha_1 \alpha_2 \alpha_3 F_{146} = 0, \\
& \alpha_1 \alpha_2 F_{127} + \alpha_2 \alpha_3 F_{154} + \alpha_3 \alpha_1 F_{163} + \alpha_1 F_{567} + \alpha_2 F_{347} + \alpha_3 F_{246} + \alpha_1 \alpha_2 \alpha_3 F_{253} = 0.
\end{aligned} \tag{3.2}$$

In particular, the $\text{SO}(1, 2)$ invariance, the M2-brane worldvolume Lorentz symmetry, removes any world-volume dependence, $D_\mu X_I = 0$ for all μ and I .

The above set of BPS equations can be regarded as the master equations since any $N = 2k$ BPS equations can be obtained by imposing k copies of distinct $(\alpha_1, \alpha_2, \alpha_3)$ choices. The corresponding $N = 2k$ BPS equations are then $\text{SO}(1, 2) \times \text{SO}(8-k) \times \text{SO}(k)$ invariant. We find for $N = 14$ and $N = 16$ the corresponding BPS equations are trivial, $F_{\mu I} = F_{IJK} = 0$. Other nontrivial cases are as follows.

3.1.1 $N = 2$ $\text{SO}(1, 2) \times \text{SO}(7)$ invariant BPS equations - octonion

With the choice of $(\alpha_1, \alpha_2, \alpha_3) = (+, +, +)$, the $N = 2$ $\text{SO}(1, 2) \times \text{SO}(7)$ invariant BPS equations (3.1), (3.2) assume a compact form:

$$F_{\mu I} = 0, \quad \mathcal{C}_{IJKL} F^{JKL} = 0, \tag{3.3}$$

where $\mathcal{C}_{IJKL}$ is a $\text{SO}(7)$ invariant four-form in eight dimensions, defined in terms of the octonionic structure constant (2.14),

$$\mathcal{C}_{ijk8} \equiv c_{ijk}, \quad \mathcal{C}_{ijkl} \equiv \frac{1}{6} \epsilon_{pqrijkl} c_{pqr} \quad \text{where} \quad 1 \leq i, j, k, l \leq 7. \tag{3.4}$$

BPS states preserving $N = 2k$ supersymmetries then satisfy k copies of the $N = 2$ BPS equations of different α choices. The corresponding $N = 2k$ BPS equations are $\text{SO}(1, 2) \times \text{SO}(k) \times \text{SO}(8-k)$ invariant, and involve k different octonionic structures.

3.1.2 $N = 4$ $\text{SO}(1, 2) \times \text{SO}(6) \times \text{SO}(2)$ invariant BPS equations - complex

The $N = 4$ $\text{SO}(1, 2) \times \text{SO}(6) \times \text{SO}(2)$ invariant BPS equations are, with $F_{\mu I} = 0$,

$$F_{IJK} \mathcal{J}^{JK} = 0, \quad F_{IJK} = (1 \otimes \mathcal{J} \otimes \mathcal{J} + \mathcal{J} \otimes 1 \otimes \mathcal{J} + \mathcal{J} \otimes \mathcal{J} \otimes 1)_{IJK}{}^{LMN} F_{LMN}, \quad (3.5)$$

where $\mathcal{J}$ is a complex structure $\mathcal{J}^2 = -1$, $\mathcal{J}^T = -\mathcal{J}$ and hence $\text{SU}(4) \times \text{SO}(2)$ invariant.

With the specific choice of α 's as $(+++), (++-)$, one gets

$$\frac{1}{2} \mathcal{J}_{IJ} \Gamma^{IJ} = \Gamma^{12} + \Gamma^{34} + \Gamma^{56} + \Gamma^{78}. \quad (3.6)$$

In terms of the corresponding holomorphic, anti-holomorphic coordinates $a, \bar{a} = 1, 2, 3, 4$ and the metric $\delta^{a\bar{a}}$, the above $N = 4$ $\text{SO}(1, 2) \times \text{SO}(6) \times \text{SO}(2)$ BPS equations (3.5) can be rewritten as

$$F_{ab}{}^b = F_{\bar{a}\bar{b}}{}^{\bar{b}} = 0, \quad F_{abc} = F_{\bar{a}\bar{b}\bar{c}} = 0. \quad (3.7)$$

Namely $F_{(1,2)}$, $F_{(2,1)}$ are primitive and $F_{(3,0)} = F_{(0,3)} = 0$.

We note that summing two $N = 2$ projection matrices generates one complex structure. Hence in general, summing $k > 2$ of $N = 2$ projection matrices will present $\binom{k}{2}$ number of complex structures to the corresponding $\text{SO}(1, 2) \times \text{SO}(8-k) \times \text{SO}(k)$ invariant BPS equations. The $\frac{1}{2}k(k-1)$ complex structures form singlets under $\text{SO}(8-k)$ and are in the adjoint representation or k -dimensional two-form representation of $\text{SO}(k)$. In fact, they correspond to the generators of $\text{SO}(k)$. Nevertheless, the corresponding $\frac{1}{2}k(k-1)$ number of complex structures are degenerate in the sense that distinct $\lceil \frac{k+1}{2} \rceil$ of them are sufficient to lead to the full $N = 2k$ BPS equations.

3.1.3 $N = 6$ $\text{SO}(1, 2) \times \text{SO}(5) \times \text{SO}(3)$ invariant BPS equations - *quaternion*

The $N = 6$ $\text{SO}(1, 2) \times \text{SO}(5) \times \text{SO}(3)$ invariant BPS equations are, with $F_{\mu I} = 0$,

$$F_{IJK} \mathcal{J}_p^{JK} = 0, \quad p = 1, 2, 3, \quad (3.8)$$

where $\mathcal{J}_1, \mathcal{J}_2, \mathcal{J}_3$ are three distinct complex structures satisfying the quaternion relations:

$$\mathcal{J}_1^2 = \mathcal{J}_2^2 = \mathcal{J}_3^2 = \mathcal{J}_1 \mathcal{J}_2 \mathcal{J}_3 = -1. \quad (3.9)$$

It is worth to note that the remaining relation of (3.5) i.e. $F_{(3,0)} = 0$ is fulfilled automatically for each complex structure.

With the specific choice of α 's as $(+++), (++-), (+-+)$, one gets

$$\begin{aligned} \frac{1}{2} \mathcal{J}_1^{IJ} \Gamma_{IJ} &= \Gamma^{12} + \Gamma^{34} + \Gamma^{56} + \Gamma^{78}, \\ \frac{1}{2} \mathcal{J}_2^{IJ} \Gamma_{IJ} &= \Gamma^{14} + \Gamma^{23} + \Gamma^{58} + \Gamma^{67}, \\ \frac{1}{2} \mathcal{J}_3^{IJ} \Gamma_{IJ} &= \Gamma^{13} + \Gamma^{42} + \Gamma^{57} + \Gamma^{86}. \end{aligned} \quad (3.10)$$

Summing three $N = 2$ projection matrices generates one quaternion structure. Hence in general, summing $k > 3$ of $N = 2$ projection matrices will present $\binom{k}{3}$ number of quaternion structures to the corresponding $\text{SO}(1, 2) \times \text{SO}(8-k) \times \text{SO}(k)$ invariant BPS equations. The $\binom{k}{3}$ quaternion structures are singlets under $\text{SO}(8-k)$ and form a k -dimensional three-form representation of $\text{SO}(k)$. Nevertheless, the corresponding $\frac{1}{6}k(k-1)(k-2)$ number of quaternion structures are degenerate in the sense that distinct $\left[\frac{k+2}{3}\right]$ of them are sufficient to give the full $N = 2k$ BPS equations.

3.1.4 $N = 8$ $\text{SO}(1, 2) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations

The $N = 8$ $\text{SO}(1, 2) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations are, with $F_{\mu I} = 0$,

$$F_{IJK} + \frac{1}{2} F_I^{LM} \mathcal{T}_{JKLM} + \frac{1}{2} F_J^{LM} \mathcal{T}_{KILM} + \frac{1}{2} F_K^{LM} \mathcal{T}_{IJLM} = 0, \quad (3.11)$$

where $\mathcal{T}_{IJKL}$ is a $\text{SO}(4) \times \text{SO}(4)$ invariant self-dual four-form. With the specific choice of α 's as $(+++), (++-), (+-+), (+--)$, one gets

$$\frac{1}{4!} \mathcal{T}_{IJKL} \Gamma^{IJKL} = \Gamma^{1234} + \Gamma^{5678}. \quad (3.12)$$

Summing four $N=2$ projection matrices generates one self-dual four-form structure. Hence in general, summing $k > 4$ of $N=2$ projection matrices will present $\binom{k}{4}$ number of self-dual four-form structures to the corresponding $\text{SO}(1,2) \times \text{SO}(8-k) \times \text{SO}(k)$ invariant BPS equations. The $\binom{k}{4}$ self-dual four-form structures are singlets under $\text{SO}(8-k)$ and form a k -dimensional four-form representation of $\text{SO}(k)$. Nevertheless, the corresponding $\frac{k!}{4!(k-4)!}$ number of self-dual four-forms are degenerate in the sense that distinct $[\frac{k+3}{4}]$ of them are sufficient to give the full $N=2k$ BPS equations.

3.1.5 $N=10$ $\text{SO}(1,2) \times \text{SO}(3) \times \text{SO}(5)$ invariant BPS equations

For $N=10$ $\text{SO}(1,2) \times \text{SO}(3) \times \text{SO}(5)$ case there seems no novel structure to appear. One economic fashion to write the $N=10$ $\text{SO}(1,2) \times \text{SO}(3) \times \text{SO}(5)$ invariant BPS equations is to employ a $\text{SO}(4) \times \text{SO}(4)$ invariant self-dual four-form and a complex structure: with $F_{\mu I} = 0$,²

$$F_{IJK} + \frac{3}{2} F_{[I}^{LM} \mathcal{T}_{JK]LM} = 0, \quad F_{IJK} \mathcal{J}^{JK} = 0. \quad (3.13)$$

The specific choice of α 's as $(+++), (++-), (+-+), (+--), (-++)$ gives

$$\frac{1}{4!} \mathcal{T}_{IJKL} \Gamma^{IJKL} = \Gamma^{1234} + \Gamma^{5678}, \quad \frac{1}{2} \mathcal{J}_{IJ} \Gamma^{IJ} = \Gamma^{18} - \Gamma^{27} + \Gamma^{36} - \Gamma^{45}. \quad (3.14)$$

3.1.6 $N=12$ $\text{SO}(1,2) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations

The $N=12$ $\text{SO}(1,2) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations are, with $F_{\mu I} = 0$,³

$$F_{IJK} \mathcal{T}_p^{JK} = 0, \quad p = 1, 2, 3, 4, 5, 6, \quad (3.15)$$

² Alternatively we can express them in terms of two sets of *either* $\text{SO}(4) \times \text{SO}(4)$ invariant self-dual four-forms one given by (3.12) the other by $\frac{1}{2}(\Gamma_{1234} + \Gamma_{5678} + \Gamma_{1256} + \Gamma_{3478} + \Gamma_{1357} + \Gamma_{2468} + \Gamma_{1467} + \Gamma_{2358})$ or quaternionic complex structures one by (3.10) and the other by $\Gamma_{14} + \Gamma_{85} + \Gamma_{76} + \Gamma_{23}$, $\Gamma_{15} + \Gamma_{48} + \Gamma_{73} + \Gamma_{62}$, $\Gamma_{18} + \Gamma_{54} + \Gamma_{72} + \Gamma_{36}$.

³ Of course, the above $N=12$ BPS equations can be obtained by imposing a pair of two distinct quaternionic BPS equations (3.8). There are $\frac{1}{2} \binom{6}{3} = 10$ such pairs and any of them leads to the same $N=12$ BPS equations. For example we may choose one quaternion structure from (3.10) and the other by $\Gamma^{12} + \Gamma^{87} + \Gamma^{56} + \Gamma^{43}$, $\Gamma^{17} + \Gamma^{28} + \Gamma^{53} + \Gamma^{64}$, $\Gamma^{18} + \Gamma^{72} + \Gamma^{54} + \Gamma^{36}$, corresponding to the α choices $(+++), (++-), (+-+)$ and $(+--), (-++), (-+-)$.

where $\mathcal{T}_p^{IJ}$'s are $\text{SO}(2) \times \text{SO}(6)$ covariant two-forms: fundamental under $\text{SO}(6)$ and singlet under $\text{SO}(2)$.

With the specific choice of α 's as $(+++),(++-),(+--),(+--),(-++),(-+-)$, one gets

$$\begin{aligned}\frac{1}{2}\mathcal{T}_1^{IJ}\Gamma_{IJ} &= \Gamma^{14} + \Gamma^{23}, & \frac{1}{2}\mathcal{T}_2^{IJ}\Gamma_{IJ} &= \Gamma^{67} + \Gamma^{58}, \\ \frac{1}{2}\mathcal{T}_3^{IJ}\Gamma_{IJ} &= \Gamma^{16} + \Gamma^{25}, & \frac{1}{2}\mathcal{T}_4^{IJ}\Gamma_{IJ} &= \Gamma^{74} + \Gamma^{83}, \\ \frac{1}{2}\mathcal{T}_5^{IJ}\Gamma_{IJ} &= \Gamma^{17} + \Gamma^{28}, & \frac{1}{2}\mathcal{T}_6^{IJ}\Gamma_{IJ} &= \Gamma^{35} + \Gamma^{46}.\end{aligned}\tag{3.16}$$

3.2 $\text{SO}(2)$ invariant BPS equations

The generic $N = 2$ projection matrix (2.21) leads to the following $N=2$ $\text{SO}(2) \times \text{SU}(4)$ invariant BPS equations which involve three free sign factors $\beta_1^2 = \beta_2^2 = \beta_3^2 = 1$:

$$\begin{aligned}F_{x1} + \beta_1 F_{y2} &= 0, & F_{x3} + \beta_2 F_{y4} &= 0, & F_{x5} + \beta_3 F_{y6} &= 0, & F_{x7} + \beta_1 \beta_2 \beta_3 F_{y8} &= 0, \\ F_{x2} - \beta_1 F_{y1} &= 0, & F_{x4} - \beta_2 F_{y3} &= 0, & F_{x6} - \beta_3 F_{y5} &= 0, & F_{x8} - \beta_1 \beta_2 \beta_3 F_{y7} &= 0,\end{aligned}\tag{3.17}$$

and

$$\begin{aligned}F_{t1} + \beta_2 F_{134} + \beta_3 F_{156} + \beta_1 \beta_2 \beta_3 F_{178} &= 0, & F_{135} - \beta_1 \beta_2 F_{245} - \beta_2 \beta_3 F_{146} - \beta_3 \beta_1 F_{236} &= 0, \\ F_{t2} + \beta_2 F_{234} + \beta_3 F_{256} + \beta_1 \beta_2 \beta_3 F_{278} &= 0, & F_{136} - \beta_1 \beta_2 F_{246} + \beta_2 \beta_3 F_{145} + \beta_3 \beta_1 F_{235} &= 0, \\ F_{t3} + \beta_1 F_{312} + \beta_3 F_{356} + \beta_1 \beta_2 \beta_3 F_{378} &= 0, & F_{137} - \beta_1 \beta_2 F_{247} - \beta_2 \beta_3 F_{238} - \beta_3 \beta_1 F_{148} &= 0, \\ F_{t4} + \beta_1 F_{412} + \beta_3 F_{456} + \beta_1 \beta_2 \beta_3 F_{478} &= 0, & F_{138} - \beta_1 \beta_2 F_{248} + \beta_2 \beta_3 F_{237} + \beta_3 \beta_1 F_{147} &= 0, \\ F_{t5} + \beta_1 F_{512} + \beta_2 F_{534} + \beta_1 \beta_2 \beta_3 F_{578} &= 0, & F_{157} - \beta_1 \beta_2 F_{168} - \beta_2 \beta_3 F_{258} - \beta_3 \beta_1 F_{267} &= 0, \\ F_{t6} + \beta_1 F_{612} + \beta_2 F_{634} + \beta_1 \beta_2 \beta_3 F_{678} &= 0, & F_{158} + \beta_1 \beta_2 F_{167} + \beta_2 \beta_3 F_{257} - \beta_3 \beta_1 F_{268} &= 0, \\ F_{t7} + \beta_1 F_{712} + \beta_2 F_{734} + \beta_3 F_{756} &= 0, & F_{357} - \beta_1 \beta_2 F_{368} - \beta_2 \beta_3 F_{467} - \beta_3 \beta_1 F_{458} &= 0, \\ F_{t8} + \beta_1 F_{812} + \beta_2 F_{834} + \beta_3 F_{856} &= 0, & F_{358} + \beta_1 \beta_2 F_{367} - \beta_2 \beta_3 F_{468} + \beta_3 \beta_1 F_{457} &= 0.\end{aligned}\tag{3.18}$$

The above set of BPS equations can be regarded as the master equations since any $N=2k$ $\text{SO}(2)^5$ invariant BPS equations corresponding to the projection matrices (2.23 - 2.27) can be obtained by imposing k copies

of distinct $(\beta_1, \beta_2, \beta_3)$ choices. We find, among them, the $N = 8$ $\text{SO}(2) \times \text{SU}(4)$ invariant projection matrix (2.27) leads to the trivial BPS configuration $F_{\mu I} = F_{IJK} = 0$. Other nontrivial cases are as follows.

3.2.1 $N = 2$ $\text{SO}(2) \times \text{SU}(4)$ invariant BPS equations

The $N = 2$ $\text{SO}(2) \times \text{SU}(4)$ invariant BPS equations corresponding to the projection matrix (2.23) or the choice $(\beta_1, \beta_2, \beta_3) = (+ + +)$ in (3.17) and (3.18) assume a compact form, up to Hermitian conjugation:

$$F_{z\bar{a}} = 0, \quad F_{ta} - iF_{ab}{}^b = 0, \quad F_{abc} = 0, \quad (3.19)$$

provided we complexify the $\text{SO}(8)$ coordinates by the complex structure $\Gamma_{12} + \Gamma_{34} + \Gamma_{56} + \Gamma_{78}$, to introduce the holomorphic and anti-holomorphic variables $a, \bar{a} = 1, 2, 3, 4$ such that the metric is $\delta_{a\bar{a}}$ and

$$\begin{aligned} D_z &= \frac{1}{\sqrt{2}}(D_x - iD_y), & D_{\bar{z}} &= \frac{1}{\sqrt{2}}(D_x + iD_y), \\ F_{za} &= \frac{1}{\sqrt{2}}(D_z X_{2a-1} - iD_z X_{2a}), & F_{z\bar{a}} &= \frac{1}{\sqrt{2}}(D_z X_{2\bar{a}-1} + iD_z X_{2\bar{a}}). \end{aligned} \quad (3.20)$$

3.2.2 $N = 4$ $\text{SO}(2) \times \text{SU}(2) \times \text{SO}(4)$ invariant BPS equations

The $N = 4$ $\text{SO}(2) \times \text{SU}(2) \times \text{SO}(4)$ invariant BPS equations corresponding to the projection matrix (2.24) are, up to Hermitian conjugation,

$$F_{z\bar{a}} = 0, \quad F_{zp} = 0, \quad F_{pab} = 0, \quad F_{tI} - iF_{Ia}{}^a = 0, \quad F_{Ipq} + \frac{1}{2}\epsilon_{pqrs}F_I{}^{rs} = 0, \quad (3.21)$$

where $I = 1, 2, \dots, 8$, $p, q, r, s = 5, 6, 7, 8$, ϵ_{pqrs} is a totally anti-symmetric tensor with $\epsilon_{5678} = 1$ and $a, b, \bar{a} = 1, 2$ such that the $\text{SO}(4) \subset \text{SO}(8)$ coordinates are complexified by the complex structure $\Gamma_{12} + \Gamma_{34}$.

3.2.3 $N = 6$ $\text{SO}(2) \times \text{SO}(2) \times \text{SU}(3)$ invariant BPS equations

The $N = 6$ $\text{SO}(2) \times \text{SO}(2) \times \text{SU}(3)$ invariant BPS equations corresponding to the projection matrix (2.25) are, up to Hermitian conjugation,

$$\begin{aligned} F_{z\bar{\omega}} &= 0, & F_{za} &= 0, & F_{z\bar{a}} &= 0, & F_{t\omega} - i\frac{1}{3}F_{\omega a}{}^a &= 0, \\ F_{ta} - iF_{a\omega\bar{\omega}} &= 0, & F_{\omega ab} &= 0, & F_{ab\bar{c}} &= 0, & F_{\omega a\bar{b}} - \frac{1}{3}(F_{\omega c}{}^c)\delta_{a\bar{b}} &= 0, \end{aligned} \quad (3.22)$$

where $a, \bar{a} = 1, 2, 3$ such that we complexify the $\text{SO}(6) \subset \text{SO}(8)$ coordinates by the complex structure $\Gamma_{34} + \Gamma_{56} + \Gamma_{87}$ and also set separately for $\text{SO}(2) \subset \text{SO}(8)$,

$$F_{z\omega} \equiv \frac{1}{\sqrt{2}}(F_{z1} - iF_{z2}), \quad F_{z\bar{\omega}} \equiv \frac{1}{\sqrt{2}}(F_{z1} + iF_{z2}). \quad (3.23)$$

3.2.4 $N = 8$ $\text{SO}(2) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations

The $N = 8$ $\text{SO}(2) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations corresponding to the projection matrix (2.26) are, up to Hermitian conjugation,

$$F_{z\bar{\omega}} = 0, \quad F_{zp} = 0, \quad F_{tI} - iF_{I\omega\bar{\omega}} = 0, \quad F_{Ipq} = 0, \quad (3.24)$$

where $I = 1, 2, \dots, 8$, $p = 3, 4, 5, 6, 7, 8$ and we complexify the $\text{SO}(2) \subset \text{SO}(8)$ coordinates by the complex structure Γ_{12} to employ (3.23).

3.3 $\text{SO}(1, 1)$ invariant BPS equations

The generic $N = 1$ projection matrix (2.28) leads to the following $N = 1$ $\text{SO}(1, 1) \times \text{SO}(7)$ invariant BPS equations which involve four free signs $\alpha_0^2 = \alpha_1^2 = \alpha_2^2 = \alpha_3^2 = 1$:

$$\begin{aligned} F_{tI} - \alpha_0 F_{xI} &= 0, \quad I = 1, 2, \dots, 8, \\ \alpha_0 F_{y1} - \alpha_1 \alpha_2 F_{278} - \alpha_2 \alpha_3 F_{548} - \alpha_3 \alpha_1 F_{638} - \alpha_1 F_{234} - \alpha_2 F_{256} - \alpha_3 F_{357} - \alpha_1 \alpha_2 \alpha_3 F_{476} &= 0, \\ \alpha_0 F_{y2} - \alpha_1 \alpha_2 F_{718} - \alpha_2 \alpha_3 F_{376} - \alpha_3 \alpha_1 F_{475} - \alpha_1 F_{143} - \alpha_2 F_{165} - \alpha_3 F_{468} - \alpha_1 \alpha_2 \alpha_3 F_{538} &= 0, \\ \alpha_0 F_{y3} - \alpha_1 \alpha_2 F_{456} - \alpha_2 \alpha_3 F_{267} - \alpha_3 \alpha_1 F_{168} - \alpha_1 F_{124} - \alpha_2 F_{478} - \alpha_3 F_{517} - \alpha_1 \alpha_2 \alpha_3 F_{258} &= 0, \\ \alpha_0 F_{y4} - \alpha_1 \alpha_2 F_{536} - \alpha_2 \alpha_3 F_{158} - \alpha_3 \alpha_1 F_{257} - \alpha_1 F_{132} - \alpha_2 F_{738} - \alpha_3 F_{628} - \alpha_1 \alpha_2 \alpha_3 F_{167} &= 0, \\ \alpha_0 F_{y5} - \alpha_1 \alpha_2 F_{346} - \alpha_2 \alpha_3 F_{418} - \alpha_3 \alpha_1 F_{427} - \alpha_1 F_{678} - \alpha_2 F_{126} - \alpha_3 F_{137} - \alpha_1 \alpha_2 \alpha_3 F_{328} &= 0, \\ \alpha_0 F_{y6} - \alpha_1 \alpha_2 F_{354} - \alpha_2 \alpha_3 F_{273} - \alpha_3 \alpha_1 F_{318} - \alpha_1 F_{758} - \alpha_2 F_{152} - \alpha_3 F_{248} - \alpha_1 \alpha_2 \alpha_3 F_{174} &= 0, \\ \alpha_0 F_{y7} - \alpha_1 \alpha_2 F_{128} - \alpha_2 \alpha_3 F_{236} - \alpha_3 \alpha_1 F_{245} - \alpha_1 F_{568} - \alpha_2 F_{348} - \alpha_3 F_{153} - \alpha_1 \alpha_2 \alpha_3 F_{146} &= 0, \\ \alpha_0 F_{y8} + \alpha_1 \alpha_2 F_{127} + \alpha_2 \alpha_3 F_{154} + \alpha_3 \alpha_1 F_{163} + \alpha_1 F_{567} + \alpha_2 F_{347} + \alpha_3 F_{246} + \alpha_1 \alpha_2 \alpha_3 F_{253} &= 0. \end{aligned} \quad (3.25)$$

The above set of BPS equations can be regarded as the master equations for generic $\text{SO}(1, 1)$ invariant BPS equations. One can classify the BPS equations according to the decomposition of the number of preserved supersymmetries as (N_+, N_-) (2.30). Among others, below we spell explicitly $(N_+, 0)$ as well as (N, N) BPS equations with $N_+ = 1, 2, \dots, 7$, $N = 1, 2, 3, 4$.

3.3.1 $(N_+, N_-) = (1, 0)$ $\text{SO}(1, 1) \times \text{SO}(7)$ invariant BPS equations - octonion

With the choice of $(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = (+ + + +)$, the $(N_+, N_-) = (1, 0)$ $\text{SO}(1, 1) \times \text{SO}(7)$ invariant BPS equations (3.25) assume a compact form:

$$F_{tI} - F_{xI} = 0, \quad F_{yI} - \frac{1}{6} C_{IJKL} F^{JKL} = 0, \quad (3.26)$$

which generalizes the $N = 2$ $\text{SO}(1, 2) \times \text{SO}(7)$ invariant BPS equations (3.3).

3.3.2 $(N_+, N_-) = (2, 0)$ $\text{SO}(1, 1) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations - complex

The $(N_+, N_-) = (2, 0)$ $\text{SO}(1, 1) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations are, with $F_{tI} - F_{xI} = 0$,

$$\mathcal{J}^{IJ} F_{yJ} + \frac{1}{2} F^I{}_{JK} \mathcal{J}^{JK} = 0, \quad F_{IJK} = (1 \otimes \mathcal{J} \otimes \mathcal{J} + \mathcal{J} \otimes 1 \otimes \mathcal{J} + \mathcal{J} \otimes \mathcal{J} \otimes 1)_{IJK}{}^{LMN} F_{LMN}, \quad (3.27)$$

which generalizes the $N = 4$ $\text{SO}(1, 2) \times \text{SO}(6) \times \text{SO}(2)$ invariant BPS equations (3.5).

3.3.3 $(N_+, N_-) = (3, 0)$ $\text{SO}(1, 1) \times \text{SO}(3) \times \text{SO}(5)$ invariant BPS equations - quaternion

The $(N_+, N_-) = (3, 0)$ $\text{SO}(1, 1) \times \text{SO}(3) \times \text{SO}(5)$ invariant BPS equations are, with $F_{tI} - F_{xI} = 0$,

$$\mathcal{J}_p^{IJ} F_{yJ} + \frac{1}{2} F^I{}_{JK} \mathcal{J}_p^{JK} = 0, \quad p = 1, 2, 3, \quad (3.28)$$

where $\mathcal{J}_1, \mathcal{J}_2, \mathcal{J}_3$ are three distinct complex structures satisfying the quaternion relations, $\mathcal{J}_1^2 = \mathcal{J}_2^2 = \mathcal{J}_3^2 = \mathcal{J}_1 \mathcal{J}_2 \mathcal{J}_3 = -1$ (3.10). It is worth to note that the remaining relation of (3.27) $F_{(3,0)} = 0$ is fulfilled automatically for each complex structure. Eq.(3.28) generalizes the $N = 6$ $\text{SO}(1, 2) \times \text{SO}(5) \times \text{SO}(3)$ invariant BPS equations (3.8).

3.3.4 $(N_+, N_-) = (4, 0)$ $\text{SO}(1, 1) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations

The $(N_+, N_-) = (4, 0)$ $\text{SO}(1, 1) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations are, with $F_{tI} - F_{xI} = 0$,

$$\mathcal{T}_{IJKL} F_y{}^L + F_{IJK} + \frac{1}{2} F_I{}^{LM} \mathcal{T}_{JKLM} + \frac{1}{2} F_J{}^{LM} \mathcal{T}_{KILM} + \frac{1}{2} F_K{}^{LM} \mathcal{T}_{IJLM} = 0, \quad (3.29)$$

where $\mathcal{T}_{IJKL}$ is a $\text{SO}(4) \times \text{SO}(4)$ invariant self-dual four-form (3.12). Eq.(3.29) generalizes the $N = 8$ $\text{SO}(1, 2) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations (3.11). Some mass deformations of the above BPS equations are studied in Ref. [23].

3.3.5 $(N_+, N_-) = (5, 0)$ $\text{SO}(1, 1) \times \text{SO}(5) \times \text{SO}(3)$ invariant BPS equations

The $(N_+, N_-) = (5, 0)$ $\text{SO}(1, 1) \times \text{SO}(5) \times \text{SO}(3)$ invariant BPS equations are, with $F_{tI} - F_{xI} = 0$,

$$\mathcal{T}_{IJKL} F_y^L + F_{IJK} + \frac{3}{2} F_I^{LM} \mathcal{T}_{JK]LM} = 0, \quad \mathcal{J}^{IJ} F_{yJ} + \frac{1}{2} F_{IJK} \mathcal{J}^{JK} = 0, \quad (3.30)$$

where $\mathcal{T}_{IJKL}$ and $\mathcal{J}^{IJ}$ are given in (3.14). Eq.(3.30) generalizes the $N = 10$ $\text{SO}(1, 2) \times \text{SO}(3) \times \text{SO}(5)$ invariant BPS equations (3.13).

3.3.6 $(N_+, N_-) = (6, 0)$ $\text{SO}(1, 1) \times \text{SO}(6) \times \text{SO}(2)$ invariant BPS equations

The $(N_+, N_-) = (6, 0)$ $\text{SO}(1, 1) \times \text{SO}(6) \times \text{SO}(2)$ invariant BPS equations are, $F_{tI} - F_{xI} = 0$,

$$\mathcal{T}_p^{IJ} F_{yJ} + \frac{1}{2} F^I_{JK} \mathcal{T}_p^{JK} = 0, \quad p = 1, 2, 3, 4, 5, 6, \quad (3.31)$$

where six of two-forms $\mathcal{T}_p$, $p = 1, 2, \dots, 6$ are given in (3.16). Eq.(3.31) generalizes the $N = 12$ $\text{SO}(1, 2) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations (3.15).

3.3.7 $(N_+, N_-) = (7, 0)$ $\text{SO}(1, 1) \times \text{SO}(7)$ invariant BPS equations

The $(N_+, N_-) = (7, 0)$ $\text{SO}(1, 1) \times \text{SO}(7)$ invariant BPS equations are, with $F_{tI} - F_{xI} = 0$,

$$\mathcal{T}_p^{IJ} F_{yJ} + \frac{1}{2} F^I_{JK} \mathcal{T}_p^{JK} = 0, \quad p = 1, 2, 3, 4, 5, 6, 7. \quad (3.32)$$

Here we have seven of two-forms, six given by (3.16) and last one by

$$\frac{1}{2} \mathcal{T}_7^{IJ} \Gamma_{IJ} = \Gamma^{13} + \Gamma^{57}. \quad (3.33)$$

They form a fundamental representation of $\text{SO}(7)$.

3.3.8 $(N_+, N_-) = (1, 1)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(6)$ invariant BPS equations

The $(N_+, N_-) = (1, 1)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(6)$ invariant BPS equations are, with $F_{tI} = F_{xI} = 0$, best expressed in complex coordinates,

$$F_{ab}{}^b = 0, \quad F_{y\bar{a}} - \frac{1}{3}\epsilon_{\bar{a}}{}^{bcd}F_{bcd} = 0. \quad (3.34)$$

3.3.9 $(N_+, N_-) = (2, 2)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(2) \times \mathbf{SO}(2) \times \mathbf{SO}(4)$ invariant BPS equations

The $(N_+, N_-) = (2, 2)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(2) \times \mathbf{SO}(2) \times \mathbf{SO}(4)$ invariant BPS equations are, with $F_{tI} = F_{xI} = 0$,

$$(3\mathcal{J}^{[IJ}\mathcal{J}^{K]L} - \mathcal{T}^{IJKL})F_{yL} + F^{IJK} + \frac{3}{2}F^{[I}{}_{LM}\mathcal{T}^{JK]LM} = 0, \quad (3.35)$$

where $\mathcal{J}^{IJ}$ is the complex structure of $\Gamma^{12} + \Gamma^{34} + \Gamma^{56} + \Gamma^{78}$ (3.6) and $\mathcal{T}^{IJKL}$ is the self-dual $\mathbf{SO}(4) \times \mathbf{SO}(4)$ invariant four-form tensor of $\Gamma^{1234} + \Gamma^{5678}$ (3.12).

3.3.10 $(N_+, N_-) = (3, 3)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(3) \times \mathbf{SO}(3) \times \mathbf{SO}(2)$ invariant BPS equations

We present the $(N_+, N_-) = (3, 3)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(3) \times \mathbf{SO}(3) \times \mathbf{SO}(2)$ invariant BPS equations with a pair of quaternion structures, one from (3.10) and the other from $\Gamma^{12} + \Gamma^{87} + \Gamma^{56} + \Gamma^{43}$, $\Gamma^{17} + \Gamma^{28} + \Gamma^{53} + \Gamma^{64}$, $\Gamma^{18} + \Gamma^{72} + \Gamma^{54} + \Gamma^{36}$. With $F_{tI} = F_{xI} = 0$ they are

$$\mathcal{J}_p^{IJ}F_{yJ} + \frac{1}{2}F^I{}_{JK}\mathcal{J}_p^{JK} = 0, \quad \hat{\mathcal{J}}_p^{IJ}F_{yJ} - \frac{1}{2}F^I{}_{JK}\hat{\mathcal{J}}_p^{JK} = 0, \quad p = 1, 2, 3. \quad (3.36)$$

3.3.11 $(N_+, N_-) = (4, 4)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(4) \times \mathbf{SO}(4)$ invariant BPS equations

The $(N_+, N_-) = (4, 4)$ $\mathbf{SO}(1, 1) \times \mathbf{SO}(4) \times \mathbf{SO}(4)$ invariant BPS equations are, with $F_{tI} = F_{xI} = 0$, in terms of the self-dual $\mathbf{SO}(4) \times \mathbf{SO}(4)$ invariant four-form tensor,

$$\mathcal{T}^{IJKL}F_{yL} + F^{IJK} = 0. \quad (3.37)$$

Especially among all the half BPS cases *i.e.* $N_+ + N_- = 8$, only the case $(N_+, N_-) = (4, 4)$ leads to the nontrivial BPS equations.

4 Discussion

In this paper we studied and identified a number of BPS equations for the multiple M2-brane theory proposed recently by Bagger and Lambert. We employed a method which had been successfully applied to several analogous problems. One first constructs the basic projection matrices for the supersymmetry parameters, and then obtain the corresponding BPS equations. Our classifications are complete for $\text{SO}(1, 2)$ as well as $\text{SO}(2)$ ⁵ invariant BPS equations, while may be not for $\text{SO}(1, 1)$ invariant cases.

The BPS equations with different types and numbers of preserved supersymmetries are derived in terms of the associated tensors which are invariant under the symmetry group of the relevant BPS equations. In particular we derived three types of half BPS equations, which we recall:

- $N = 8$ $\text{SO}(1, 2) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations (3.11)

$$F_{\mu I} = 0, \quad F_{IJK} + \frac{1}{2}F_I^{LM}\mathcal{T}_{JKLM} + \frac{1}{2}F_J^{LM}\mathcal{T}_{KILM} + \frac{1}{2}F_K^{LM}\mathcal{T}_{IJLM} = 0. \quad (4.1)$$

- $N = 8$ $\text{SO}(2) \times \text{SO}(2) \times \text{SO}(6)$ invariant BPS equations (3.24)

$$F_{z\bar{\omega}} = 0, \quad F_{zp} = 0, \quad F_{tI} - iF_{I\omega\bar{\omega}} = 0, \quad F_{Ipq} = 0, \quad (4.2)$$

where $I = 1, 2, \dots, 8$, $p = 3, 4, 5, 6, 7, 8$, and $\omega, \bar{\omega}$ are complex coordinates for $\text{SO}(2) \subset \text{SO}(8)$.

- $(N_+, N_-) = (4, 4)$ $\text{SO}(1, 1) \times \text{SO}(4) \times \text{SO}(4)$ invariant BPS equations (3.37)

$$F_{tI} = F_{xI} = 0, \quad \mathcal{T}^{IJKL}F_{yL} + F^{IJK} = 0. \quad (4.3)$$

The BPS equations for different number of supersymmetries exhibit the division algebra structures: octonion, quaternion or complex. Let us take the Lorentz invariant type as examples. For the least supersymmetric configurations preserving 1/8 supersymmetries, the relevant symmetry is $\text{SO}(1, 2) \times \text{SO}(7)$ and the BPS equations can be elegantly written in terms of the invariant four-form which has close relation to octonions. For 1/4-BPS equations the symmetry is $\text{SO}(1, 2) \times \text{SO}(6) \times \text{SO}(2)$ and a complex structure appears. We next have 3/8 $\text{SO}(1, 2) \times \text{SO}(5) \times \text{SO}(3)$ invariant BPS equations, which are naturally best expressed in terms of quaternions or hyper-Kähler structure. In addition, for 1/2-BPS equations we have the $\text{SO}(4) \times \text{SO}(4)$ invariant self-dual four-form structure. We have also identified the exotic classes with more

than 1/2 supersymmetry. Apparently the governing symmetries include more than one hyper-Kähler structures, but we have not been able to express the BPS equations in a succinct way. The true mathematical identity of such systems certainly deserves more careful study.

The explicit solutions of the BPS equations will give the spectrum of supersymmetric solitons in Bagger-Lambert theory. It is natural to ask the $\mathcal{M}$ -theory interpretation of such objects. The real scalar fields X^I describe the locations of M2-branes in the transverse $\mathbb{R}^8$. The spatial dependence of X^I thus informs us on the shape of M2-branes, or how they are embedded in the transverse $\mathbb{R}^8$. Eq.(3.17) and the subsequent analysis clearly suggest that the M2-brane worldvolume should occupy holomorphic curves, which is natural for supersymmetry. Likewise, time-dependence of the scalar field obviously implies that there is momentum along the particular direction. The three-algebra terms F_{IJK} describe the truly $\mathcal{M}$ -theoretic phenomena: polarization of multiple M2-branes into M5-branes. Generically the BPS equations are given as various combinations of such basic building blocks, and more detailed descriptions with explicit solutions will be reported in a separate publication.

Acknowledgments

We wish to thank Bum-Hoon Lee for discussions and encouragement. This work is supported by the Center for Quantum Spacetime of Sogang University with grant number R11 - 2005 - 021. NK is partly supported by Korea Research Foundation Grant, No. KRF-2007-331-C00072. The research of JHP is supported in part by the Korea Science and Engineering Foundation grant funded by the Korea government (R01-2007-000-20062-0).

A Gamma matrices and octonions

The eleven-dimensional 32×32 gamma matrices Γ^M , $M = \mu, I, \mu = t, x, y, I = 1, 2, \dots, 8$ in the Bagger-Lambert theory naturally decompose into two parts: $\text{SO}(1, 2)$ the M2-brane worldvolume and $\text{SO}(8)$ the transverse space,

$$\Gamma^t = \epsilon \otimes \gamma_{(9)}, \quad \Gamma^x = \sigma_1 \otimes \gamma_{(9)}, \quad \Gamma^y = \sigma_3 \otimes \gamma_{(9)}, \quad \Gamma^I = 1 \otimes \gamma^I, \quad I = 1, 2, \dots, 8. \quad (\text{A.1})$$

Here γ^I 's are the 16×16 gamma matrices in the eight-dimensional Euclidean space and $\gamma_{(9)} \equiv \gamma_{12 \dots 8}$. Clearly the $\text{SO}(1, 2)$ projection constraint (2.4) coincides with that of $\text{SO}(8)$,

$$\Gamma^{txy} = 1 \otimes \gamma_{(9)}. \quad (\text{A.2})$$

This is consistent with the fact that the product of all the eleven-dimensional gamma matrices leads to the identity $\Gamma^{txy123 \dots 8} = 1$.

Now we recall the seven quantities $\mathcal{E}_i$, $i = 1, 2, 3 \dots, 7$ (2.13). In the above choice of gamma matrices we have

$$\mathcal{E}_i = 1 \otimes E_i, \quad \mathcal{P} = 1 \otimes P, \quad (\text{A.3})$$

where as in (2.13)

$$\begin{aligned} E_1 &= \gamma_{8127} P, & E_2 &= \gamma_{8163} P, & E_3 &= \gamma_{8246} P, & E_4 &= \gamma_{8347} P, \\ E_5 &= \gamma_{8567} P, & E_6 &= \gamma_{8253} P, & E_7 &= \gamma_{8154} P, & P &= \frac{1}{2}(1 + \gamma_{(9)}). \end{aligned} \quad (\text{A.4})$$

The subscript spatial indices of the gamma matrices are organized such that the three indices after the common 8 are identical to those of the totally anti-symmetric octonionic structure constants (2.14). It is straightforward to see that E_i forms a representation of the “square” of the octonions on the eight-dimensional chiral space,

$$E_i E_j = \delta_{ij} P + c_{ijk}^2 E_k, \quad E_i \equiv e_i \otimes e_i. \quad (\text{A.5})$$

Since they commute each other, they form a maximal set of the mutually commuting traceless symmetric and real matrices of the definite chirality $\gamma_{(9)} E_i = E_i$. In fact, one can construct a $\text{SO}(8)$ symmetric

and real gamma matrix representation which makes all E_i 's be simultaneously diagonal, utilizing the octonionic structure constants:

$$\gamma_I = \begin{pmatrix} 0 & \rho_I \\ (\rho_I)^T & 0 \end{pmatrix}, \quad \rho_I(\rho_J)^T + \rho_J(\rho_I)^T = 2\delta_{IJ}, \quad \gamma_{(9)} = \gamma_{12345678} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (\text{A.6})$$

Here ρ_I , $I = 1, 2, \dots, 8$ are 8×8 real matrices given by⁴

$$\rho_i = -(\rho_i)^T = \begin{pmatrix} c_i & -n_i \\ (n_i)^T & 0 \end{pmatrix}, \quad i = 1, 2, \dots, 7, \quad \rho_8 = 1, \quad (\text{A.7})$$

and c_i is a 7×7 real matrix whose j, k component is nothing but the octonionic structure constant c_{ijk} (2.14), while n_i is a seven-dimensional unit vector of which the j th component is defined to be δ_i^j .

In the above choice of Majorana gamma matrix representation, all the E_i 's and P are diagonal,

$$\begin{aligned} E_1 &= \text{diag}(+1, +1, -1, -1, -1, -1, +1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ E_2 &= \text{diag}(+1, -1, +1, -1, -1, +1, -1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ E_3 &= \text{diag}(-1, +1, -1, +1, -1, +1, -1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ E_4 &= \text{diag}(-1, -1, +1, +1, -1, -1, +1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ E_5 &= \text{diag}(-1, -1, -1, -1, +1, +1, +1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ E_6 &= \text{diag}(-1, +1, +1, -1, +1, -1, -1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ E_7 &= \text{diag}(+1, -1, -1, +1, +1, -1, -1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \\ P &= \text{diag}(+1, +1, +1, +1, +1, +1, +1, +1, 0, 0, 0, 0, 0, 0, 0, 0), \end{aligned} \quad (\text{A.8})$$

and the $\text{SO}(8)$ triality among $\mathbf{8}_v$, $\mathbf{8}_+$, $\mathbf{8}_-$ is apparent as the $\mathbf{8}_v$ generators decompose into the $\mathbf{8}_+$ and $\mathbf{8}_-$ generators,

$$\gamma_{IJ} = \begin{pmatrix} \rho_{[I}\rho_{J]}^T & 0 \\ 0 & \rho_{[I}^T\rho_{J]} \end{pmatrix}. \quad (\text{A.9})$$

⁴In particular, ρ_i , $1 \leq i \leq 7$ correspond to the Majorana gamma matrices in Euclidean seven dimensions $\rho_i\rho_j + \rho_j\rho_i = -2\delta_{ij}$.

With the identity $e_8 \equiv 1$, the octonion algebra now spells completely:

$$e_I e_J = (\rho_I)_{JK} e_K, \quad I, J, K = 1, 2, \dots, 8. \quad (\text{A.10})$$

Finally let us consider a self-dual four-form and contract it with the $\text{SO}(8)$ gamma matrices Γ^{IJKL} , such as $\Upsilon_4 \mathcal{P}$ in (2.10). Clearly utilizing the $\text{SO}(8)$ triality, one can diagonalize $\Upsilon_4 \mathcal{P}$ to express it as a linear combination of $\mathcal{E}_i$'s. This shows that the canonical form of a self-dual four-form in eight dimensions indeed takes the form (2.12): namely the non-vanishing independent components are only those seven which are contracted to $\mathcal{E}_i$'s.

B $\text{SO}(2)$ invariant projection matrix

Here we derive the most general form of the 32×32 projection matrices Ω which are invariant under the Cartan subalgebra $\text{SO}(2)^5$ of $\text{SO}(10)$, satisfying in addition to the conditions (2.8),

$$[\Gamma^{xy}, \Omega] = 0, \quad [\Gamma^{12}, \Omega] = 0, \quad [\Gamma^{34}, \Omega] = 0, \quad [\Gamma^{56}, \Omega] = 0, \quad [\Gamma^{78}, \Omega] = 0. \quad (\text{B.1})$$

As (2.17), they assume the general form:

$$\Omega = [c + \Gamma^{xy}(a_1 \Gamma^{12} + a_2 \Gamma^{34} + a_3 \Gamma^{56} + a_4 \Gamma^{78}) + b_1 \Gamma^{1234} + b_2 \Gamma^{1256} + b_3 \Gamma^{1278}] \mathcal{P}, \quad (\text{B.2})$$

where $c, a_1, \dots, b_3$ are eight *a priori* unknown real constants which must be determined by requiring the remaining condition $\Omega^2 = \Omega$. In particular the number of the preserved supersymmetries is related to the constant c by

$$N = \text{Tr} \Omega = 16c. \quad (\text{B.3})$$

It is convenient to reparameterize the four constants a_1, a_2, a_3, a_4 by four other constants e_1, e_2, e_3, e_4

$$\begin{aligned} e_1 &= 2(a_1 + a_2 + a_3 + a_4), & e_2 &= 2(a_1 + a_2 - a_3 - a_4), \\ e_3 &= 2(a_1 - a_2 + a_3 - a_4), & e_4 &= 2(-a_1 + a_2 + a_3 - a_4), \end{aligned} \quad (\text{B.4})$$

and the other four constants c, b_1, b_2, b_3 by another set of four constants f_1, f_2, f_3, f_4

$$\begin{aligned} f_1 &= 2c - 1 - 2b_1 - 2b_2 - 2b_3, & f_2 &= 2c - 1 - 2b_1 + 2b_2 + 2b_3, \\ f_3 &= 2c - 1 + 2b_1 - 2b_2 + 2b_3, & f_4 &= 2c - 1 + 2b_1 + 2b_2 - 2b_3. \end{aligned} \quad (\text{B.5})$$

It follows that

$$\begin{aligned}
a_1 &= \frac{1}{8}(e_1 + e_2 + e_3 - e_4), & a_2 &= \frac{1}{8}(e_1 + e_2 - e_3 + e_4), \\
a_3 &= \frac{1}{8}(e_1 - e_2 + e_3 + e_4), & a_4 &= \frac{1}{8}(e_1 - e_2 - e_3 - e_4), \\
b_1 &= \frac{1}{8}(-f_1 - f_2 + f_3 + f_4), & b_2 &= \frac{1}{8}(-f_1 + f_2 - f_3 + f_4), \\
b_3 &= \frac{1}{8}(-f_1 + f_2 + f_3 - f_4), & c &= \frac{1}{8}(f_1 + f_2 + f_3 + f_4 + 4).
\end{aligned} \tag{B.6}$$

Straightforward calculation shows that $\Omega^2 = \Omega$ is equivalent for each $a = 1, 2, 3, 4$ to

$$f_a e_a = 0, \quad e_a^2 = (1 + f_a)(1 - f_a) \quad \text{not } a \text{ sum.} \tag{B.7}$$

Hence for each a we have four possible solutions:

$$e_a = 0, f_a = +1; \quad e_a = 0, f_a = -1; \quad e_a = +1, f_a = 0; \quad e_a = -1, f_a = 0. \tag{B.8}$$

Consequently from (B.3) and (B.6), the possible values of c are $0, \frac{1}{8}, \frac{2}{8}, \frac{3}{8}, \frac{4}{8}, \frac{5}{8}, \frac{6}{8}, \frac{7}{8}, 1$, so that the number of the preserved supersymmetries N is an even number between zero and sixteen. The basic building blocks of all the possible projection matrices are those of $N = 2$ given by

$$\begin{aligned}
\Omega &= \frac{1}{8} [1 + \Gamma^{xy} (\beta_1 \Gamma^{12} + \beta_2 \Gamma^{34} + \beta_3 \Gamma^{56} + \beta_1 \beta_2 \beta_3 \Gamma^{78}) - \beta_1 \beta_2 \Gamma^{1234} - \beta_3 \beta_1 \Gamma^{1256} - \beta_2 \beta_3 \Gamma^{1278}] \mathcal{P} \\
&= \frac{1}{8} (1 + \beta_1 \Gamma^{xy12}) (1 + \beta_2 \Gamma^{xy34}) (1 + \beta_3 \Gamma^{xy56}) \mathcal{P},
\end{aligned} \tag{B.9}$$

where $\beta_1, \beta_2, \beta_3$ are three independent signs,

$$\beta_1^2 = \beta_2^2 = \beta_3^2 = 1. \tag{B.10}$$

There are eight possible $N = 2$ projection matrices which are orthogonal to each other. By summing k of them, all the other generic projection matrices preserving $N = 2k$ supersymmetries can be obtained.

References

- [1] J. Bagger and N. Lambert, Phys. Rev. D **75** (2007) 045020 [arXiv:hep-th/0611108];
J. Bagger and N. Lambert, Phys. Rev. D **77** (2008) 065008 [arXiv:0711.0955 [hep-th]];
J. Bagger and N. Lambert, JHEP **0802** (2008) 105 [arXiv:0712.3738 [hep-th]].
- [2] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, arXiv:0803.3242 [hep-th].
- [3] A. Basu and J. A. Harvey, Nucl. Phys. B **713** (2005) 136 [arXiv:hep-th/0412310].
- [4] A. Gustavsson, arXiv:0709.1260 [hep-th].
- [5] O. J. Ganor, Phys. Rev. D **75** (2007) 025002 [arXiv:hep-th/0609107].
- [6] P. M. Ho and Y. Matsuo, Gen. Rel. Grav. **39** (2007) 913 [arXiv:hep-th/0701130].
- [7] N. B. Copland, arXiv:0707.1317 [hep-th].
- [8] B. Chen, W. He, J. B. Wu and L. Zhang, JHEP **0708** (2007) 067 [arXiv:0707.3978 [hep-th]].
- [9] O. A. P. Mac Conamhna, arXiv:0708.2568 [hep-th].
- [10] D. S. Berman, Phys. Rept. **456** (2008) 89 [arXiv:0710.1707 [hep-th]].
- [11] B. Chen, Eur. Phys. J. C **54** (2008) 489 [arXiv:0710.2593 [hep-th]].
- [12] I. A. Bandos and J. A. de Azcarraga, JHEP **0804** (2008) 069 [arXiv:0802.2890 [hep-th]].
- [13] A. Gustavsson, JHEP **0804** (2008) 083 [arXiv:0802.3456 [hep-th]].
- [14] S. Mukhi and C. Papageorgakis, arXiv:0803.3218 [hep-th].
- [15] D. S. Berman, L. C. Tadowski and D. C. Thompson, arXiv:0803.3611 [hep-th].
- [16] M. Van Raamsdonk, arXiv:0803.3803 [hep-th].
- [17] A. Morozov, arXiv:0804.0913 [hep-th].
- [18] N. Lambert and D. Tong, arXiv:0804.1114 [hep-th].

- [19] U. Gran, B. E. W. Nilsson and C. Petersson, arXiv:0804.1784 [hep-th].
- [20] P. M. Ho, R. C. Hou and Y. Matsuo, arXiv:0804.2110 [hep-th].
- [21] J. Gomis, A. J. Salim and F. Passerini, arXiv:0804.2186 [hep-th].
- [22] E. A. Bergshoeff, M. de Roo and O. Hohm, arXiv:0804.2201 [hep-th].
- [23] K. Hosomichi, K. M. Lee and S. Lee, arXiv:0804.2519 [hep-th].
- [24] G. Papadopoulos, arXiv:0804.2662 [hep-th].
- [25] J. P. Gauntlett and J. B. Gutowski, arXiv:0804.3078 [hep-th].
- [26] H. Shimada, arXiv:0804.3236 [hep-th].
- [27] G. Papadopoulos, arXiv:0804.3567 [hep-th].
- [28] P. M. Ho and Y. Matsuo, arXiv:0804.3629 [hep-th].
- [29] J. Gomis, G. Milanesi and J. G. Russo, arXiv:0805.1012 [hep-th].
- [30] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni and H. Verlinde, arXiv:0805.1087 [hep-th].
- [31] P. M. Ho, Y. Imamura and Y. Matsuo, arXiv:0805.1202 [hep-th].
- [32] A. Morozov, arXiv:0805.1703 [hep-th].
- [33] Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, arXiv:0805.1895 [hep-th].
- [34] H. Fuji, S. Terashima and M. Yamazaki, arXiv:0805.1997 [hep-th].
- [35] P. M. Ho, Y. Imamura, Y. Matsuo and S. Shiba, arXiv:0805.2898 [hep-th].
- [36] T. Banks, N. Seiberg and S. H. Shenker, Nucl. Phys. B **490** (1997) 91 [arXiv:hep-th/9612157].
- [37] I. Jeon, J. Kim, N. Kim, S.-W. Kim, B.-H. Lee and J.-H. Park, *in preparation*.
- [38] C. Krishnan and C. Maccaferri, arXiv:0805.3125 [hep-th].
- [39] J.-H. Park and D. Tsimpis, Nucl. Phys. B **776** (2007) 405 [arXiv:hep-th/0610159].

- [40] D. Bak, K. M. Lee and J.-H. Park, Phys. Rev. D **66** (2002) 025021 [arXiv:hep-th/0204221].
- [41] D. E. Berenstein, J. M. Maldacena and H. S. Nastase, JHEP **0204** (2002) 013 [arXiv:hep-th/0202021].
- [42] J.-H. Park, JHEP **0210** (2002) 032 [arXiv:hep-th/0208161].
- [43] N. Kim and J.-H. Park, Phys. Rev. D **66** (2002) 106007 [arXiv:hep-th/0207061].
- [44] J. C. Baez, “The Octonions,” arXiv:math.ra/0105155.